

Meditation and Yoga Practices as Potential Adjunctive Treatment of SARS-CoV-2 Infection and COVID-19: A Brief Overview of Key Subjects

William Bushell, PhD,^{1,2} Ryan Castle, BS,² Michelle A. Williams, ScD,³ Kimberly C. Brouwer, PhD,⁴ Rudolph E. Tanzi, PhD,⁵ Deepak Chopra, MD,^{2,6} and Paul J. Mills, PhD⁷

Editor's Note: As an acute condition quickly associated with multiple chronic susceptibilities, COVID-19 has rekindled interest in, and controversy about, the potential role of the host in disease processes. While hundreds of millions of research dollars have been funneled into drug and vaccine solutions that target the external agent, integrative practitioners tuned to enhancing immunity faced a familiar mostly unfunded task. First, go to school on the virus. Then draw from the global array of natural therapies and practices with host-enhancing or anti-viral capabilities to suggest integrative treatment strategies. The near null-set of conventional treatment options propels this investigation. In this paper, researchers from the Massachusetts Institute of Technology, University of California-San Diego, Chopra Library for Integrative Studies, and Harvard University share one such exploration. Their conclusion, that “certain meditation, yoga asana (postures), and pranayama (breathing) practices may possibly be effective adjunctive means of treating and/or preventing SARS-CoV-2 infection” underscores the importance of this rekindling. At JACM, we are pleased to have the opportunity to publish this work. We hope that it might help diminish in medicine and health the polarization that, like so much in the broader culture, seems to be an obstacle to healing. —John Weeks, Editor-in-Chief, JACM

Potential Adjunctive Therapies for SARS-CoV-2 Infection and COVID-19 Disease

THERE IS AN URGENT need to identify strategies to help prevent and treat SARS-CoV-2 infection and COVID-19 disease.^{1–4} Included among the intensive search for novel and effective therapeutic approaches are considerations of those therapies derived from integrative and complementary medicine,^{5–8} including from traditional whole medicine systems such as Chinese Medicine and Ayurvedic medicine.^{6–8} In this article, the authors present a selective narrative review of the literature with a primary focus on certain complementary practices of seated meditation, yoga asanas (postures), and pranayama (breathing) with relevance to immune function for consideration for SARS-CoV-2 infection and COVID-19 treatment and/or prevention.

The authors propose that certain complementary practices may be helpful *adjunctive* means of treating and/or

preventing SARS-CoV-2 infection^{1,9} and helping to reduce severity of COVID-19 disease, including its collateral effects and sequelae. Despite being speculative for the present situation, there is a body of literature relevant to the anti-stress and anti-inflammatory effects of certain seated meditation, yoga asanas, and pranayama practices. Such studies include demonstrating promising immune effects relevant to improving lung health^{10,11} and reducing viral susceptibility and improving acute respiratory infections.^{12–14} The potential benefits of these practices extend to broader neuroimmune systems, an advantage when dealing with a systemically dysregulating disease such as SARS-CoV-2.¹⁵ Such complementary practices have been found to act as a regulating influence on a number of key inflammatory functions that SARS-CoV-2 disrupts.¹⁶

The use of certain complementary practices as potentially effective adjunctive means of treating and/or preventing SARS-CoV-2 infection must be put to rigorous scientific

¹Biophysical/Medical Anthropology, Massachusetts Institute of Technology, Cambridge, MA, USA.

²Chopra Library for Integrative Studies, Whole Health Institute, Bentonville, AK, USA.

³Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA.

⁴Division of Global Health, Infectious Disease Epidemiology, Department of Family Medicine and Public Health, University of California, San Diego, La Jolla, CA, USA.

⁵Department of Neurology, Harvard University, Cambridge, MA, USA.

⁶Department of Family Medicine and Public Health, University of California, San Diego, La Jolla, CA, USA.

⁷Department of Family Medicine and Public Health, Center of Excellence for Integrative Health, Institute for Public Health, University of California, San Diego, La Jolla, CA, USA.

investigation. During this time, however, when there are numerous exploratory pathways being urgently probed,^{1,2,6,8} it is important to look to “evidence-based principles that emphasize the practical application of the findings of the best available current research” (see “implementation of evidence-based interventions” defined and advocated by NIH/CDC).¹⁷ With this consideration in mind, best available research reasonably advocates that the utilization of certain forms of some complementary practices be examined as potential adjunctive interventions to help prevent and treat SARS-CoV-2 infection and potential future pandemics.

Theoretical frameworks as models have been put forth seeking to explain how complementary practices such as seated meditation, yoga asanas, and pranayama work, which have examined multiple psychologic, emotional, epigenetic, neural, and behavioral processes, as well as shifts in self-awareness.^{18–24} An in-depth model proposed by Holzel et al., for example, focuses on mindfulness meditation practice-induced changes in attention regulation, body awareness, emotion regulation, and perspective of self, citing supportive underlying neuroplastic changes.¹⁸ Gard et al. examine the effects of yoga practices on cognition, emotional, and behavioral regulation, and autonomic output under stress with an emphasis on interoceptive processes and self-regulation.¹⁹

The current state of the rapidly evolving field of Integrative Medicine research intersects with the intensive search for successful treatments for this pandemic viral infection. When considering complementary practices, the National Academy of Science’s Institute of Medicine stated that the field of “behavioral medicine” should be considered in the evolving standard model of fundamental life science research and should be advocated on the front line of serious medical and health-optimizing treatment and intervention.²⁵ With this in mind, in 2009, one of the authors (W.B.), with the collaboration of Neil Theise MD (then of the Albert Einstein College of Medicine), began to review the scattered and disparately emerging body of research on the inflammatory cytokine-based pathogenesis of many virulent infectious diseases (bacterial, viral, other) and “cytokine storms,”²⁶ following the pioneering work of Clark et al.²⁷ and others. This was connected with research into the potential anti-inflammatory, antistress, and anti-infectious properties of meditation and yoga practices, thereby beginning the development of a scientific model in behavioral health.^{26,28} This developing scholarly territory needs to be fully explored with scientific rigor and open minds due to both its inherent value and the urgency of the present situation.^{29,30}

Anti-Inflammatory Effects Associated with Meditation and Yoga Practices

An extended runaway hyperinflammatory host response to SARS-CoV-2 infection is the primary pathway to COVID-19 disease morbidity and mortality.³¹ Spread of the virus through the body leads to widespread and intensive activation of the inflammatory defenses, although originally intended to combat the pathogen, but instead resulting in widespread tissue damage to the host, to acute respiratory distress syndrome (ARDS); the virus itself plays a much lesser role in the tissue damage that can ultimately produce

extreme critical disease states (pulmonary aspiration, septic shock), and potentially death.^{15,32}

Beginning in the mid-1990s and developing up to the present, increases in scientific knowledge about the immune system and the inflammatory dimension of it led to the recognition that many infectious diseases caused their damage to a large extent through the host’s own inflammatory responses.²⁷ Around the same time, research by Kevin Tracey and colleagues³³ (and see Bushell, Olivo, Theise, 2009)³⁴ led to the discovery that inflammation could be brought under control through stimulation of the vagal nerve complex, a major component of the central and peripheral nervous system, which also controls responses to psychosocial stress, and stimulation of which can reverse the “fight or flight” response through replacing it with the “relaxation response,”³⁵ in simplified schematic terms.

Starting in the early 2000s, Bushell and colleagues proposed that intensive forms of meditation and yoga could potentially be effective as adjunctive interventions against serious forms of infectious diseases, including malaria, HIV/AIDS, and SARS, among others. These authors were at the time demonstrating an appreciation of the key significance of the common denominator of the inflammatory-driven pathogenesis of the spectrum of such virulent infectious diseases and the potentially important relevance of the antistress and anti-inflammatory properties of these behavioral health practices for possible forms of adjunctive preventative or treatment therapies.^{27,36–38}

It had also been found that meditation, as well as yoga practices, is capable of significantly increasing vagal tone and therefore could not only be effective against psychologic stress-based issues, including trauma,^{35,36} but inflammatory-based diseases as well.^{16,41–46} Included in this consideration are studies documenting effects of a diverse range of meditative and yoga practices to promote anti-inflammatory and typically reduce proinflammatory cytokine activity, including natural killer (NK) cell activity and NK and T cell cytokine production,^{45–50} which has been a topic of two systematic reviews.^{51,52} For example, Bower et al.⁴⁹ showed that 6 weeks of 20-min daily mindfulness meditation resulted in a significant downregulation of a 19-transcript composite of proinflammatory genes and a significant reduction of activity of the proinflammatory transcription factor nuclear factor kappa B (NF- κ B). In parallel, they observed an increase of anti-inflammatory glucocorticoid receptor activity and increased type I interferon signaling, a transcription factor linked to SARS-CoV-2 treatment.

Other studies show decreases in the circulating levels of the proinflammatory cytokine interleukin (IL)-12 and increases of the anti-inflammatory cytokine IL-10.^{47,48,50} In specific patient groups or obese individuals, other studies document reductions in circulating levels of C-reactive protein (CRP) and IL-6,^{49,50,53} although these findings were not replicated by some investigators.⁴⁸ Some of the cytokine-related effects of seated meditation, breathing practices, and yoga asana practices are related to a reduction in sympathetic nervous system activation, as evidenced by parallel reductions in levels of the catecholamines epinephrine and norepinephrine.^{47,52,54} Studies also suggest differential effects of meditation practices on immune cell subsets, including NK and T cells, of relevance to the innate and adaptive immune systems.^{55,56}

The benefits of yoga therapies on the health of the innate immune system have also been demonstrated in a study showing that 90 min of yogic asana stretching is able to increase expression of the two important antimicrobial peptides, β -defensin and HBD-2 expression.⁵⁷ Importantly, with regard to COVID-19, both of these antimicrobial peptides are expressed in respiratory epithelial cells.⁵⁸ Yoga therapy has also been shown to downregulate the cytokine receptors tumor necrosis factor (TNF)-RII and IL-1RA, and stress-related CRP.⁵⁹ Epel et al. showed that combined mantra-based meditation and yoga asana practices regulated levels of the proinflammatory cytokine TNF- α and metabolism of the Alzheimer's disease-associated amyloid- β protein,⁶⁰ and also reported reduced expression of proinflammatory genes in association with a meditation practice.

Table 1 presents cytokine effect sizes from practices of meditation, yoga asana, and pranayama, reviewed in this article, paired with findings to date from COVID-19 studies reporting cytokine values. Figure 1 presents effect sizes from Table 1. It is evident that effect sizes are larger with COVID-19. The significance of the presentation, however, is to show the relevant overlap and general directions of effects on cytokines and that these practices can provide systemic regulatory effects on a range of cytokines. It is important to note that the reviewed studies were on more short-term practitioners of the modalities where such cytokine data are available. There is a fairly large literature, although comparing the effects of longer term compared with more novice practitioners, that shows significantly larger effects with longer practice in domains, for example, of autonomic, neurohormonal, and brain functioning.⁶¹⁻⁶³ Needed are studies of immune function in longer term practitioners directly compared with novice practitioners, although some studies have examined immune-related profiles in longer term meditators compared with nonpractitioner controls.⁶⁴⁻⁶⁶

Furthermore, according to a small but relevant body of research, certain meditation and yoga practices bring another set of key neuroendocrine factors into the picture, in particular melatonin, which in addition to its systemic

health-promoting properties has beneficial effects on mood, affect, emotion, and mental state and functioning.^{26,27,67,68} Studies provide insight into the epigenetic and microbiome pathways of complementary practices relevant to inflammatory processes.^{24,64-66,69}

Meditation and Yoga May Potentially Enhance the Activity of Melatonin, One of the Most Beneficial Pleiotropic Natural Substances Known

The functions of melatonin are extensive. In the words of leading melatonin researcher Russell Reiter of the University of Texas, writing in the life sciences journal, *Physiology*, and referring to thousands of published scientific studies⁷⁰: “Melatonin has a bewildering array of functions and employs a variety of means to carry them out...These actions likely impact every cell in every organism throughout the plant and animal kingdoms. Its discovered actions are uniformly beneficial, although not all the specific mechanisms have been described. In addition to its multiple positive physiologic actions, melatonin has an uncommonly high safety profile.” As cited in Bushell,⁶⁸ these include a wide array of anti-inflammatory, antioxidant, anti-infectious immune-enhancing actions on most of the cells and organs of the body that have been investigated, including the brain, heart, visceral organs, bone, muscle, and skin. In addition, there are beneficial effects on the mitochondria and nucleus of cells, including DNA repair. Melatonin has also been demonstrated to possess significant antiviral properties, including relevance to respiratory diseases.^{71,72}

Of particular relevance, researchers have specifically identified melatonin as a substance that should be investigated for possible treatment against COVID-19.^{73,74} Furthermore, melatonin is protective against ventilator-induced lung injury in animal models.^{75,76} Studies in animals and humans have found beneficial effects of melatonin against serious viral infections, including herpes, HIV/AIDS, influenza A-H1N1, West Nile virus, coxsackievirus (the virus that

TABLE 1. CYTOKINE EFFECT SIZES (COHEN *D*) FROM COMPLEMENTARY STUDIES PAIRED WITH FINDINGS FROM COVID-19 STUDIES REPORTING CYTOKINE VALUES

Study	COVID-19 d	Median d	Marker	Median d	Meditation d	Study
Chen et al. ³	-1.651	-1.651	IFN- λ	-0.783	-1.235	Carlson et al. ⁴⁰
Qin et al. ²⁹	0.493	0.493	IL-10	0.118	-0.330	Carlson et al. ⁴²
Chen et al. ³	-1.578	-1.578	IL-10	-0.375	-0.073	Carlson et al. ⁴⁰
			IL-4	-0.889	0.309	Witek-Janusek et al. ⁴¹
Qin et al. ²⁹	2.911	2.722	IL-6	-0.386	-0.375	Carlson et al. ⁴²
Wang et al. ⁴	2.533				-1.014	Carlson et al. ^{40,2007}
					-1.007	Carlson et al. ⁴²
					-0.459	Witek-Janusek et al. ⁴¹
					-0.571	Witek-Janusek et al. ⁴¹
					0.582	Cahn et al. ⁴⁴
					-0.941176471	Shete et al. ⁵⁵
					-0.2	Bower et al. ⁴⁵
Qin et al. ²⁹	-1.452	-1.452	IL-8		0.472	Witek-Janusek et al. ⁴¹
Qin et al. ²⁹	-3.077	-3.077	NK activity	0.472	0.472	Carlson et al. ⁴⁰
Qin et al. ²⁹	3.008	3.008	TNF- α	-1.064	-1.064	Epel et al. ⁵⁶
					-1.407	Shete et al. ⁵⁵
					-1.043	

IFN- λ , interferon- λ ; IL, interleukin; NK, natural killer; TNF- α , tumor necrosis factor alpha.




FIG. 1. Graphics of effect sizes (Cohen d) presented in Table 1. It is evident that effect sizes are larger with COVID-19. The significance of the graphics is the relevant overlap and general direction of effects on cytokines and that these practices can provide broad systemic regulatory effects on a range of cytokines. Note too that the reviewed studies were from more short-term practitioners of the modalities, where such cytokine data are available. There is a fairly large literature comparing the effects of longer term with more novice practitioners, which shows significantly larger effects with longer practice in domains of autonomic, neurohormonal, and brain functioning. Needed are studies of immune function in longer term practitioners compared with novices. The median effect sizes of complementary practices tracked here include small ($d \sim 0.2$), moderate ($d \sim 0.4$), and large ($d \sim 0.80$).

causes Venezuelan equine encephalomyelitis [VEE], Semliki Forest virus, and encephalomyocarditis virus.^{73,77–86} Melatonin has also been hypothesized to contribute to heretofore unexplained seasonality of certain infectious diseases.⁸ Mechanisms through which melatonin exerts its protective and antiviral effects include protective antioxidant and free radical-scavenging properties (e.g., HIV/AIDS, other retroviruses),^{17,88} including with respect to the potential development of septic shock (e.g., VEE; some forms of ARDS of viral origin)^{71,89}; immunomodulatory effects, which can include initial acute upregulation of inflammatory responses followed by well-timed downregulation, including through key influences on gene expression of NF- κ B and specific inflammatory cytokine-regulating genes (e.g., respiratory syncytial virus; viral encephalitis)^{77,80}; and through melatonin's recently discovered modulation of autophagic and apoptotic processes (e.g., coxsackievirus-B3, some forms of herpes virus disease).^{78,81}

While not an area of intensive research, a small yet compelling body of studies show that regular practice of meditation and/or yoga is associated with elevations in melatonin.^{87,90–95} Studies report that while during the practices of seated meditation and pranayama, levels of melatonin can decrease,^{91,96} longer term practice leads to elevations in levels.^{90,92,96} Solberg et al., for example, reported that more experienced meditators had 150% higher resting levels of melatonin (4.9 pg/mL)

compared with an age-, gender-, and physical fitness level-matched reference group (3.1 pg/mL).⁸⁷

Melatonin appears to possess the capacity to selectively up- and downregulate its antiviral/immune-enhancing and its anti-inflammatory effects according to the state of the milieu or environment in which it is present.^{78,97–100} In other words, melatonin biochemically “senses” the inflammatory and/or viral (or more generally infectious) nature of its *milieu interieur*, and then appears to generate the necessary effects to fit the situation, from its wide-ranging pleiotropic repertoire of potential effects. This is a growing field that needs further development. Whether or not it can do so successfully in the case of COVID-19 is a matter for further research,^{73,74} but presently there is research being conducted by institutions worldwide, which seek to find medicines and/or medicine regimens that possess simultaneous capacities for antiviral and anti-inflammatory actions, in line with such principles.^{73,101,102}

Meditation and Yoga for Well-Being During the Social Distancing Intervention for Ending the COVID-19 Global Pandemic: A Program That Goes Beyond Homeostasis

Relevant to the COVID-19 pandemic are the documented effects of some complementary practices, even potentially for novice practitioners, on reducing the severity of

psychosocial stressors, including social isolation. While it is recognized that certain levels of stress can be beneficial for the development of resilience,¹⁰³ the emerging stress landscape of the COVID-19 pandemic is extreme, involving not only threats to health and physical survival, but also to economic survival, housing, adequate nutrition, personal relationships, the very fabric of society and “normal life” itself, ranging from the local to global scale. The prospect of being a patient in an overwhelmed health system is among the most extreme stressors possible to experience, owing to the heightened sense of vulnerability.

These kinds of extreme chronic stressors are known to be capable of leading to an overwhelmed immune system, compromising the ability to mount an effective immune response to infectious challenge and leaving the individual susceptible to severe medical consequences.^{104,105} Scientific research shows that psychosocial stress can not only substantially reduce immune defenses against infectious challenges but also overstimulate host inflammatory responses to such a level that it can lead to tissue damage and even death.^{24,26,106,107} In addition, studies show in the elderly, a population of particular susceptibility to SARS-CoV-2, that stressors such as caregiving, chronic stress, bereavement stress, and social isolation are associated with increased inflammation, including elevated circulating cytokine levels.^{108–113}

Epidemiologic and historical epidemiologic evidence, from the 1918–19 influenza pandemic to more recent epidemics and pandemics involving SARS and H1N1, demonstrates that social distancing can serve a vital function^{114,115} in restricting further transmission of viral and other pathogens. Nevertheless, it is clear that such social distancing can also lead to intensified social isolation stress,^{116,117} which is in fact utilized as a standard immunity-compromising challenge in animal studies of viral infection and immune response. In addition to the social isolation stress, other pandemic-associated stressors, as listed above, can amount in many cases to pandemic-based PTSD.^{118,119} Because of the likely ongoing nature of the pandemic, it would perhaps make sense in this context to speak of *Ongoing* Traumatic Stress Disorder (OTSD), rather than *Post* Traumatic Stress Disorder. Bruce McEwen and colleagues developed the concepts of allostatic load and overload as quantifiable constructs for ongoing states of chronic stress—as is implied in the label OTSD—and underscored the complex pathophysiologic synergisms between psychosocial stressors and inflammation, including through epigenetic pathways, and have advocated the use of meditation techniques as treatments.^{120,121}

Germane to these current pandemic-associated psychosocial stressors, there is evidence that diverse meditation and yoga practices provide beneficial effects on mental health. For example, systematic reviews of the effects of a variety of meditation techniques on depressed mood, anxiety, and quality of life show promising findings in diverse populations, including clinical populations.^{122–124} Systematic and more qualitative reviews examining the effects of yoga practices on these same endpoints also report evidence of beneficial effects, although this literature is somewhat less consistent.^{125–127}

Regarding different forms of meditation practice, given the current social stressors, the authors posit that loving kindness and compassion meditation practices could possi-

bly be utilized to not only counteract the inherent negatives of the developing situation—from the perspectives of individual physical and emotional health—but socially as well.^{128–130} Certain forms of stress can synergize to produce greater negative consequences, but the converse may also be true that such forms of loving kindness/compassion meditation lead to a “broadening and building” of social connectedness through the generation of positive psychologic states,^{130,131} which are further associated with enhanced vagal tone¹³² and even reduced viral replication and function (e.g., in HIV-infected women),¹³³ thereby leading to a range of associated positive outcomes.

Recent articles in the scientific¹³⁴ and popular press¹³⁵ show that the COVID-19 pandemic is beginning to reveal social isolation-based negative effects on mental health, including depression and anxiety, which typically exert subsequent negative health outcomes. These considerations may be particularly relevant to the elderly who are more prone to both social isolation and its adverse psychologic effects, as well as to higher levels of chronic inflammation (recently dubbed “inflammaging”).¹³⁶

Mediation/Yoga-Based Behavioral Health Interventions for Disease Survivors and Their Progeny: The Important Role of the Lingering Effects of Infection and the “Inflammatory Load”

It is crucial to include in this brief overview a larger perspective on the potentially devastating effects of the pandemic relating to the risk of overwhelming the health care system. Modeling and projections of international health care systems show that while highly developed nations are able to absorb the extreme patient load from SARS-CoV-2 without disrupting their core functions, they will not be able to do so in the face of a sustained pandemic.¹³⁷ The strain on the system is exacerbated by the lengthy and intensive care required to combat the cytokine storms.³² Here the authors consider the potential and likely *aftereffects* of infection, from both individual and societal perspectives. A particular key term is that of inflammatory load which the life science biologist and historical epidemiologist, Caleb E Finch, developed as a major operational construct in all fields of human biologic science.¹³⁸ Inflammatory load in individuals and populations represents chronic elevated levels of inflammatory substances in the blood and tissues due to lingering subclinical levels of infectious pathogens, bacterial, viral, and otherwise. According to this body of research, the persistence of pathogens in environments contributes to increased risk and occurrence of most chronic diseases through transmission of subclinical infection, including cardiovascular, neurodegenerative, and other major forms of chronic inflammation-based diseases.

While in contemporary times a higher burden of chronic systemic inflammation has resulted from noninfectious environmental factors, including pollution,^{139,140} infectious pandemics and large-scale epidemics increase infectious cohort-wide inflammatory damage in populations, as Finch and colleagues demonstrated for the 1918 influenza pandemic and other widespread infectious disease situations.¹³⁸ As they conclude in a key article, “a ‘cohort morbidity phenotype’ represents inflammatory processes that persist from early age into adult life”.¹³⁸ These contribute to

increased risk and incidence of cardiovascular diseases, neurodegenerative diseases such as Alzheimer's, and a range of other diseases that are, in large part, pathogenically based on extended durations of chronically elevated levels of inflammatory substances in the body.^{141,142}

According to the above-cited work on the 1918 influenza pandemic, such conditions resulting from heightened inflammation also significantly affect future generations, as Finch and colleagues¹³⁸ reported, that those born to mothers infected during the pandemic had heightened vulnerability to cardiovascular disease (males particularly) and growth retardation, and that "prenatal exposure to even uncomplicated maternal influenza may have lasting consequences later in life".¹⁴³ This research is to be further considered in the context of psychosocial stress-associated influences that may extend over generations, via epigenetic and possibly other mechanisms—and especially with respect to the pathophysiological connections between psychosocial stress and inflammation.^{144,145}

Conclusion and Further Recommendations

The authors find themselves in the midst of a pandemic, the extent of which is not yet clear, and it is a time to enlist all manner of assistance in resisting and combating the multiple threats associated with it. As they have reviewed, there is evidence of stress and inflammation modulation, and also preliminary evidence for possible forms of immune system enhancement, accompanying the practice of certain forms of meditation, yoga, and pranayama, along with potential implications for counteracting some forms of infectious challenges. Their apparent association with enhanced melatonin activity also implies another pathway by which the beneficial properties of these modalities may be effective, especially with respect to aspects of viral challenges of various forms.

The points discussed in this article suggest potential initial short-term acute interventional and long-term benefits focusing on these practices as well as in the reduction of future sequelae to pandemic-associated negative inflammatory and psychosocial stress factors.

The authors readily acknowledge that in the context of the SARS-CoV-2 pandemic, the ideas put forth in this article must be put to further rigorous scientific investigation. During this time of urgency, however, when numerous novel approaches are being explored for their potential therapeutic efficacy, based on the existing evidence base for certain complementary practices as here briefly reviewed, the authors advocate for such consideration and further research.

Authors' Contributions

All authors contributed to this work.

Author Disclosure Statement

W.B. and R.C. are part of the Chopra Research Library. P.J.M. is director of research for the Chopra Foundation. D.C. is the founder of the Chopra Foundation.

Funding Information

No funding was received for this article.

References

1. Murthy S, Gomersall CD, Fowler RA. Care for critically ill patients with COVID-19. *JAMA* 2020;323:1499–1500.
2. Baden LR, Rubin EJ. Covid-19—The search for effective therapy. *N Engl J Med* 2020;382:1787–1799.
3. Chen G, Wu D, Guo W, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest* May 1 2020;130(5):2620–2629.
4. Wang F, Nie J, Wang H, et al. Characteristics of Peripheral Lymphocyte Subset Alteration in COVID-19 Pneumonia. *J Infect Dis* May 11 2020;221:1762–1769.
5. Alschuler L, Weil A, Horwitz R, et al. Integrative considerations during the COVID-19 pandemic. *Explore (NY)* 2020; DOI: 10.1016/j.explore.2020.03.007.
6. Chan KW, Wong VT, Tang SCW. COVID-19: An update on the epidemiological, clinical, preventive and therapeutic evidence and guidelines of integrative chinese-western medicine for the management of 2019 novel coronavirus disease. *Am J Chin Med* 2020;48:737–762.
7. Yang G, Zhang H, Yang Y. Challenges and countermeasures of integrative cancer therapy in the epidemic of COVID-19. *Integr Cancer Ther* 2020;19:1534735420912811.
8. Tillu G, Chaturvedi S, Chopra A, Patwardhan B. Public health approach of ayurveda and yoga for COVID-19 prophylaxis. *J Altern Complement Med* 2020;26:360–364.
9. Arentz M, Yim E, Klaff L, et al. Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington State. *JAMA* 2020; DOI: 10.1001/jama.2020.4326.
10. Paudyal P, Jones C, Grindey C, et al. Meditation for asthma: Systematic review and meta-analysis. *J Asthma* 2018;55:771–778.
11. Leung RW, McKeough ZJ, Peters MJ, Alison JA. Short-form Sun-style t'ai chi as an exercise training modality in people with COPD. *Eur Respir J* 2013;41:1051–1057.
12. Obasi CN, Brown R, Ewers T, et al. Advantage of meditation over exercise in reducing cold and flu illness is related to improved function and quality of life. *Influenza Other Respir Viruses* 2013;7:938–944.
13. Barrett B, Hayney MS, Muller D, et al. Meditation or exercise for preventing acute respiratory infection (MEPARI-2): A randomized controlled trial. *PLoS One* 2018;13:e0197778.
14. Morgan N, Irwin MR, Chung M, Wang C. The effects of mind-body therapies on the immune system: Meta-analysis. *PLoS One* 2014;9:e100903.
15. Fu Y, Cheng Y, Wu Y. Understanding SARS-CoV-2-mediated inflammatory responses: From mechanisms to potential therapeutic tools. *Virology* 2020 [Epub ahead of print]; DOI: 10.1007/s12250-020-00207-4.
16. Bower JE, Irwin MR. Mind-body therapies and control of inflammatory biology: A descriptive review. *Brain Behav Immun* 2016;51:1–11.
17. NIH/CDC. Global Health Research Topics. Implementation of evidence based interventions. Online document at: <https://www.fic.nih.gov/ResearchTopics>, accessed June 8, 2020.
18. Holzel BK, Lazar SW, Gard T, et al. How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspect Psychol Sci* 2011;6:537–559.
19. Gard T, Noggle JJ, Park CL, et al. Potential self-regulatory mechanisms of yoga for psychological health. *Front Hum Neurosci* 2014;8:770.

20. Khanna S, Greeson JM. A narrative review of yoga and mindfulness as complementary therapies for addiction. *Complement Ther Med* 2013;21:244–252.
21. Davidson RJ, Kabat-Zinn J, Schumacher J, et al. Alterations in brain and immune function produced by mindfulness meditation. *Psychosom Med* 2003;65:564–570.
22. Jacobs TL, Epel ES, Lin J, et al. Intensive meditation training, immune cell telomerase activity, and psychological mediators. *Psychoneuroendocrinology* 2011;36:664–681.
23. Bhasin MK, Dusek JA, Chang BH, et al. Relaxation response induces temporal transcriptome changes in energy metabolism, insulin secretion and inflammatory pathways. *PLoS One* 2013;8:e62817.
24. Househam AM, Peterson CT, Mills PJ, Chopra D. The effects of stress and meditation on the immune system, human microbiota, and epigenetics. *Adv Mind Body Med* 2017;31:10–25.
25. Smedley BD, Syme SL; Committee on Capitalizing on Social S; Behavioral Research to Improve the Public's H. Promoting health: Intervention strategies from social and behavioral research. *Am J Health Promot* 2001;15:149–166.
26. Bushell WC, Theise ND. Toward a unified field of study: Longevity, regeneration, and protection of health through meditation and related practices. *Ann N Y Acad Sci* 2009;1172:5–19.
27. Clark IA, Alleva LM, Mills AC, Cowden WB. Pathogenesis of malaria and clinically similar conditions. *Clin Microbiol Rev* 2004;17:509–539, table of contents.
28. Bushell WC, Spector NH, Theise ND. From the global to the local: Possible pathways for the transduction of Indo-Sino-Tibetan cognitive-behavioral practices into site-specific, tissue-regenerative effects. *Ann N Y Acad Sci* 2009;1172:74–87.
29. Swanson DR, Smalheiser NR, Bookstein A. Information discovery from complimentary literatures: A stimulus to scientific discovery. *Artif Intell* 1997;91:183–203.
30. Swanson DR, Smalheiser NR. Undiscovered public knowledge: A ten-year update. In: *KDD-96 Proceedings. Association for the Advancement of Artificial Intelligence*, 1996.
31. Mehta P, McAuley DF, Brown M, et al. COVID-19: Consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395:1033–1034.
32. Qin C, Zhou L, Hu Z, et al. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clin Infect Dis* 2020; DOI: 10.1093/cid/ciaa248.
33. Pavlov VA, Tracey KJ. The cholinergic anti-inflammatory pathway. *Brain Behav Immun* 2005;19:493–499.
34. Bushell WC, Olivo EL, Theise ND. Longevity, Regeneration, and Optimal Health. NYAS. Online document at: <https://www.nyas.org/annals/longevity-regeneration-and-optimal-health/>, accessed June 8, 2020.
35. Jacobs GD. The physiology of mind-body interactions: The stress response and the relaxation response. *J Altern Complement Med* 2001;7(Suppl. 1):S83–S92.
36. Kuntsevich V, Bushell WC, Theise ND. Mechanisms of yogic practices in health, aging, and disease. *Mt Sinai J Med* 2010;77:559–569.
37. Awandare GA, Goka B, Boeuf P, et al. Increased levels of inflammatory mediators in children with severe *Plasmodium falciparum* malaria with respiratory distress. *J Infect Dis* 2006;194:1438–1446.
38. Okabayashi T, Kariwa H, Yokota S, et al. Cytokine regulation in SARS coronavirus infection compared to other respiratory virus infections. *J Med Virol* 2006;78:417–424.
39. Boyd JE, Lanius RA, McKinnon MC. Mindfulness-based treatments for posttraumatic stress disorder: A review of the treatment literature and neurobiological evidence. *J Psychiatry Neurosci* 2018;43:7–25.
40. Nidich S, Mills PJ, Rainforth M, et al. Non-trauma-focused meditation versus exposure therapy in veterans with post-traumatic stress disorder: A randomised controlled trial. *Lancet Psychiatry* 2018;5:975–986.
41. Breit S, Kupferberg A, Rogler G, Hasler G. Vagus nerve as modulator of the brain-gut axis in psychiatric and inflammatory disorders. *Front Psychiatry* 2018;9:44.
42. Falkenberg RI, Eising C, Peters ML. Yoga and immune system functioning: A systematic review of randomized controlled trials. *J Behav Med* 2018;41:467–482.
43. Sullivan MB, Erb M, Schmalzl L, et al. Yoga therapy and polyvagal theory: The convergence of traditional wisdom and contemporary neuroscience for self-regulation and resilience. *Front Hum Neurosci* 2018;12:67.
44. Carlson LE, Specia M, Faris P, Patel KD. One year pre-post intervention follow-up of psychological, immune, endocrine and blood pressure outcomes of mindfulness-based stress reduction (MBSR) in breast and prostate cancer outpatients. *Brain Behav Immun* 2007;21:1038–1049.
45. Witek-Janusek L, Albuquerque K, Chroniak KR, et al. Effect of mindfulness based stress reduction on immune function, quality of life and coping in women newly diagnosed with early stage breast cancer. *Brain Behav Immun* 2008;22:969–981.
46. Carlson LE, Specia M, Patel KD, Goodey E. Mindfulness-based stress reduction in relation to quality of life, mood, symptoms of stress, and immune parameters in breast and prostate cancer outpatients. *Psychosom Med* 2003;65:571–581.
47. Jang JH, Park HY, Lee US, et al. Effects of mind-body training on cytokines and their interactions with catecholamines. *Psychiatry Investig* 2017;14:483–490.
48. Cahn BR, Goodman MS, Peterson CT, et al. Yoga, meditation and mind-body health: Increased BDNF, cortisol awakening response, and altered inflammatory marker expression after a 3-month yoga and meditation retreat. *Front Hum Neurosci* 2017;11:315.
49. Bower JE, Crosswell AD, Stanton AL, et al. Mindfulness meditation for younger breast cancer survivors: A randomized controlled trial. *Cancer* 2015;121:1231–1240.
50. Jedel S, Hoffman A, Merriman P, et al. A randomized controlled trial of mindfulness-based stress reduction to prevent flare-up in patients with inactive ulcerative colitis. *Digestion* 2014;89:142–155.
51. Black DS, Slavich GM. Mindfulness meditation and the immune system: A systematic review of randomized controlled trials. *Ann N Y Acad Sci* 2016;1373:13–24.
52. Pascoe MC, Thompson DR, Ski CF. Yoga, mindfulness-based stress reduction and stress-related physiological measures: A meta-analysis. *Psychoneuroendocrinology* 2017;86:152–168.
53. Malarkey WB, Jarjoura D, Klatt M. Workplace based mindfulness practice and inflammation: A randomized trial. *Brain Behav Immun* 2013;27:145–154.
54. Kox M, van Eijk LT, Zwaag J, et al. Voluntary activation of the sympathetic nervous system and attenuation of the

- innate immune response in humans. *Proc Natl Acad Sci U S A* 2014;111:7379–7384.
55. Infante JR, Peran F, Rayo JI, et al. Levels of immune cells in transcendental meditation practitioners. *Int J Yoga* 2014;7:147–151.
 56. Vera FM, Manzanique JM, Rodriguez FM, et al. Acute effects on the counts of innate and adaptive immune response cells after 1 month of taoist qigong practice. *Int J Behav Med* 2016;23:198–203.
 57. Eda N, Shimizu K, Suzuki S, et al. Effects of yoga exercise on salivary beta-defensin 2. *Eur J Appl Physiol* 2013;113:2621–2627.
 58. Kim J, Yang YL, Jang SH, Jang YS. Human beta-defensin 2 plays a regulatory role in innate antiviral immunity and is capable of potentiating the induction of antigen-specific immunity. *Virology* 2018;15:124.
 59. Shete SU, Verma A, Kulkarni DD, Bhogal RS. Effect of yoga training on inflammatory cytokines and C-reactive protein in employees of small-scale industries. *J Educ Health Promot* 2017;6:76.
 60. Epel ES, Puterman E, Lin J, et al. Meditation and vacation effects have an impact on disease-associated molecular phenotypes. *Transl Psychiatry* 2016;6:e880.
 61. Brand S, Holsboer-Trachsler E, Naranjo JR, Schmidt S. Influence of mindfulness practice on cortisol and sleep in long-term and short-term meditators. *Neuropsychobiology* 2012;65:109–118.
 62. Wagner Robb S, Haslam A, Wirth MD, et al. Relationship between meditation and waking salivary cortisol secretion among long-term MBSR instructors. *Complement Med Res* 2019;26:101–109.
 63. Kral TRA, Schuyler BS, Mumford JA, et al. Impact of short- and long-term mindfulness meditation training on amygdala reactivity to emotional stimuli. *Neuroimage* 2018;181:301–313.
 64. Kaliman P, Alvarez-Lopez MJ, Cosin-Tomas M, et al. Rapid changes in histone deacetylases and inflammatory gene expression in expert meditators. *Psychoneuroendocrinology* 2014;40:96–107.
 65. Chaix R, Fagny M, Cosin-Tomas M, et al. Differential DNA methylation in experienced meditators after an intensive day of mindfulness-based practice: Implications for immune-related pathways. *Brain Behav Immun* 2020; 84:36–44.
 66. Chaix R, Alvarez-Lopez MJ, Fagny M, et al. Epigenetic clock analysis in long-term meditators. *Psychoneuroendocrinology* 2017;85:210–214.
 67. Bushell WC. From molecular biology to anti-aging cognitive-behavioral practices: The pioneering research of Walter Pierpaoli on the pineal and bone marrow foreshadows the contemporary revolution in stem cell and regenerative biology. *Ann N Y Acad Sci* 2005;1057: 28–49.
 68. Bushell WC. Serum factor that restores youthful function to apparently senescent stem cells is identified by recently developed expert decision tree-guided bioinformatics program. In: *Control and Regulation of Stem Cells*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2008.
 69. Kaliman P. Epigenetics and meditation. *Curr Opin Psychol* 2019;28:76–80.
 70. Reiter RJ, Tan DX, Galano A. Melatonin: Exceeding expectations. *Physiology (Bethesda)* 2014;29:325–333.
 71. Srinivasan V, Mohamed M, Kato H. Melatonin in bacterial and viral infections with focus on sepsis: A review. *Recent Pat Endocr Metab Immune Drug Discov* 2012;6: 30–39.
 72. Habtemariam S, Daglia M, Sureda A, et al. Melatonin and respiratory diseases: A review. *Curr Top Med Chem* 2017; 17:467–488.
 73. Zhou Y, Hou Y, Shen J, et al. Network-based drug repurposing for novel coronavirus 2019-nCoV/SARS-CoV-2. *Cell Discov* 2020;6:14.
 74. Zhang R, Wang X, Ni L, et al. COVID-19: Melatonin as a potential adjuvant treatment. *Life Sci* 2020;250:117583.
 75. Pedreira PR, Garcia-Prieto E, Parra D, et al. Effects of melatonin in an experimental model of ventilator-induced lung injury. *Am J Physiol Lung Cell Mol Physiol* 2008; 295:L820–L827.
 76. Wu GC, Peng CK, Liao WI, et al. Melatonin receptor agonist protects against acute lung injury induced by ventilator through up-regulation of IL-10 production. *Respir Res* 2020;21:65.
 77. Bonilla E, Valero N, Chacin-Bonilla L, Medina-Leendertz S. Melatonin and viral infections. *J Pineal Res* 2004;36: 73–79.
 78. Boga JA, Coto-Montes A, Rosales-Corral SA, et al. Beneficial actions of melatonin in the management of viral infections: A new use for this “molecular handyman”? *Rev Med Virol* 2012;22:323–338.
 79. Laliena A, San Miguel B, Crespo I, et al. Melatonin attenuates inflammation and promotes regeneration in rabbits with fulminant hepatitis of viral origin. *J Pineal Res* 2012;53:270–278.
 80. Huang SH, Cao XJ, Wei W. Melatonin decreases TLR3-mediated inflammatory factor expression via inhibition of NF-kappa B activation in respiratory syncytial virus-infected RAW264.7 macrophages. *J Pineal Res* 2008;45: 93–100.
 81. Sang Y, Gu X, Pan L, et al. Melatonin ameliorates coxsackievirus B3-induced myocarditis by regulating apoptosis and autophagy. *Front Pharmacol* 2018;9: 1384.
 82. Montiel M, Bonilla E, Valero N, et al. Melatonin decreases brain apoptosis, oxidative stress, and CD200 expression and increased survival rate in mice infected by Venezuelan equine encephalitis virus. *Antivir Chem Chemother* 2015;24:99–108.
 83. Ben-Nathan D, Maestroni GJ, Lustig S, Conti A. Protective effects of melatonin in mice infected with encephalitis viruses. *Arch Virol* 1995;140:223–230.
 84. Masters A, Pandi-Perumal SR, Seixas A, et al. Melatonin, the hormone of darkness: From sleep promotion to ebola treatment. *Brain Disord Ther* 2014;4:1000151.
 85. Anderson G, Maes M, Markus RP, Rodriguez M. Ebola virus: Melatonin as a readily available treatment option. *J Med Virol* 2015;87:537–543.
 86. Bonilla E, Valero-Fuenmayor N, Pons H, Chacin-Bonilla L. Melatonin protects mice infected with Venezuelan equine encephalomyelitis virus. *Cell Mol Life Sci* 1997; 53:430–434.
 87. CDC. Emerging Infectious Diseases. Online document at: <https://wwwnc.cdc.gov/eid/article/7/3/01-7301> article, accessed June 8, 2020.
 88. Zhang Z, Araghi-Niknam M, Liang B, et al. Prevention of immune dysfunction and vitamin E loss by dehydro-

- epiandrosterone and melatonin supplementation during murine retrovirus infection. *Immunology* 1999;96:291–297.
89. Valero N, Espina LM, Mosquera J. Melatonin decreases nitric oxide production, inducible nitric oxide synthase expression and lipid peroxidation induced by Venezuelan encephalitis equine virus in neuroblastoma cell cultures. *Neurochem Res* 2006;31:925–932.
 90. Liou CH, Hsieh CW, Hsieh CH, et al. Detection of nighttime melatonin level in Chinese Original Quiet Sitting. *J Formos Med Assoc* 2010;109:694–701.
 91. Martarelli D, Cocchioni M, Scuri S, Pompei P. Diaphragmatic breathing reduces exercise-induced oxidative stress. *Evid Based Complement Alternat Med* 2011;2011:932430.
 92. Harinath K, Malhotra AS, Pal K, et al. Effects of Hatha yoga and Omkar meditation on cardiorespiratory performance, psychologic profile, and melatonin secretion. *J Altern Complement Med* 2004;10:261–268.
 93. Tooley GA, Armstrong SM, Norman TR, Sali A. Acute increases in night-time plasma melatonin levels following a period of meditation. *Biol Psychol* 2000;53:69–78.
 94. Massion AO, Teas J, Hebert JR, et al. Meditation, melatonin and breast/prostate cancer: Hypothesis and preliminary data. *Med Hypotheses* 1995;44:39–46.
 95. Devi SK, Chansauria JP, Udupa KN. Mental depression and kundalini yoga. *Anc Sci Life* 1986;6:112–118.
 96. Solberg EE, Holen A, Ekeberg O, et al. The effects of long meditation on plasma melatonin and blood serotonin. *Med Sci Monit* 2004;10:CR96–CR101.
 97. Carrillo-Vico A, Lardone PJ, Alvarez-Sanchez N, et al. Melatonin: Buffering the immune system. *Int J Mol Sci* 2013;14:8638–8683.
 98. Markus RP, Fernandes PA, Kinker GS, et al. Immune-pineal axis—Acute inflammatory responses coordinate melatonin synthesis by pinealocytes and phagocytes. *Br J Pharmacol* 2018;175:3239–3250.
 99. Hardeland R. Melatonin and inflammation—Story of a double-edged blade. *J Pineal Res* 2018;65:e12525.
 100. Hardeland R. Aging, melatonin, and the pro- and anti-inflammatory networks. *Int J Mol Sci* 2019;20:1223.
 101. Stebbing J, Phelan A, Griffin I, et al. COVID-19: Combining antiviral and anti-inflammatory treatments. *Lancet Infect Dis* 2020;20:400–402.
 102. Vautrin A, Manchon L, Garcel A, et al. Both anti-inflammatory and antiviral properties of novel drug candidate ABX464 are mediated by modulation of RNA splicing. *Sci Rep* 2019;9:792.
 103. Ashokan A, Sivasubramanian M, Mitra R. Seeding stress resilience through inoculation. *Neural Plast* 2016;2016:4928081.
 104. Segerstrom SC, Miller GE. Psychological stress and the human immune system: A meta-analytic study of 30 years of inquiry. *Psychol Bull* 2004;130:601–630.
 105. Song H, Fall K, Fang F, et al. Stress related disorders and subsequent risk of life threatening infections: Population based sibling controlled cohort study. *BMJ* 2019;367:15784.
 106. Steptoe A, Hamer M, Chida Y. The effects of acute psychological stress on circulating inflammatory factors in humans: A review and meta-analysis. *Brain Behav Immun* 2007;21:901–912.
 107. Rohleder N. Acute and chronic stress induced changes in sensitivity of peripheral inflammatory pathways to the signals of multiple stress systems—2011 Curt Richter Award Winner. *Psychoneuroendocrinology* 2012;37:307–316.
 108. von Kanel R, Dimsdale JE, Mills PJ, et al. Effect of Alzheimer caregiving stress and age on frailty markers interleukin-6, C-reactive protein, and D-dimer. *J Gerontol A Biol Sci Med Sci* 2006;61:963–969.
 109. von Kanel R, Mills PJ, Mausbach BT, et al. Effect of Alzheimer caregiving on circulating levels of C-reactive protein and other biomarkers relevant to cardiovascular disease risk: A longitudinal study. *Gerontology* 2012;58:354–365.
 110. Butcher SK, Lord JM. Stress responses and innate immunity: Aging as a contributory factor. *Aging Cell* 2004;3:151–160.
 111. McDade TW, Hawkey LC, Cacioppo JT. Psychosocial and behavioral predictors of inflammation in middle-aged and older adults: The Chicago health, aging, and social relations study. *Psychosom Med* 2006;68:376–381.
 112. Chirinos DA, Ong JC, Garcini LM, et al. Bereavement, self-reported sleep disturbances, and inflammation: Results from project HEART. *Psychosom Med* 2019;81:67–73.
 113. Yang YC, Li T, Frenk SM. Social network ties and inflammation in U.S. adults with cancer. *Biodemography Soc Biol* 2014;60:21–37.
 114. Paynter S, Ware RS, Shanks GD. Host and environmental factors reducing mortality during the 1918–1919 influenza pandemic. *Epidemiol Infect* 2011;139:1425–1430.
 115. Reperant LA, Osterhaus A. AIDS, Avian flu, SARS, MERS, Ebola, Zika... what next? *Vaccine* 2017;35(35 Pt A):4470–4474.
 116. Cacioppo JT, Cacioppo S, Capitanio JP, Cole SW. The neuroendocrinology of social isolation. *Annu Rev Psychol* 2015;66:733–767.
 117. Mumtaz F, Khan MI, Zubair M, Dehpour AR. Neurobiology and consequences of social isolation stress in animal model—A comprehensive review. *Biomed Pharmacother* 2018;105:1205–1222.
 118. Sprang G, Silman M. Posttraumatic stress disorder in parents and youth after health-related disasters. *Disaster Med Public Health Prep* 2013;7:105–110.
 119. Xu J, Zheng Y, Wang M, et al. Predictors of symptoms of posttraumatic stress in Chinese university students during the 2009 H1N1 influenza pandemic. *Med Sci Monit* 2011;17:PH60–PH64.
 120. Fava GA, McEwen BS, Guidi J, et al. Clinical characterization of allostatic overload. *Psychoneuroendocrinology* 2019;108:94–101.
 121. McEwen BS. In pursuit of resilience: Stress, epigenetics, and brain plasticity. *Ann N Y Acad Sci* 2016;1373:56–64.
 122. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: A systematic review and meta-analysis. *JAMA Intern Med* 2014;174:357–368.
 123. Hilton L, Hempel S, Ewing BA, et al. Mindfulness meditation for chronic pain: Systematic review and meta-analysis. *Ann Behav Med* 2017;51:199–213.
 124. Edwards MK, Loprinzi PD. Comparative effects of meditation and exercise on physical and psychosocial health outcomes: A review of randomized controlled trials. *Postgrad Med* 2018;130:222–228.
 125. Cramer H, Anheyer D, Lauche R, Dobos G. A systematic review of yoga for major depressive disorder. *J Affect Disord* 2017;213:70–77.

126. Cramer H, Lauche R, Anheyer D, et al. Yoga for anxiety: A systematic review and meta-analysis of randomized controlled trials. *Depress Anxiety* 2018;35:830–843.
127. Grossl EJ, Chopra D, Mills PJ. An overview of yoga research for health and well-being. *J Yoga Phys Ther* 2015;5:210–214.
128. Fredrickson BL, Boulton AJ, Firestine AM, et al. Positive emotion correlates of meditation practice: A comparison of mindfulness meditation and loving-kindness meditation. *Mindfulness (NY)* 2017;8:1623–1633.
129. Hofmann SG, Grossman P, Hinton DE. Loving-kindness and compassion meditation: Potential for psychological interventions. *Clin Psychol Rev* 2011;31:1126–1132.
130. Hutcherson CA, Seppala EM, Gross JJ. Loving-kindness meditation increases social connectedness. *Emotion* 2008;8:720–724.
131. Fredrickson BL, Joiner T. Reflections on positive emotions and upward spirals. *Perspect Psychol Sci* 2018;13:194–199.
132. Kok BE, Coffey KA, Cohn MA, et al. How positive emotions build physical health: Perceived positive social connections account for the upward spiral between positive emotions and vagal tone. *Psychol Sci* 2013;24:1123–1132.
133. Wilson TE, Weedon J, Cohen MH, et al. Positive affect and its association with viral control among women with HIV infection. *Health Psychol* 2017;36:91–100.
134. Mannix R, Lee LK, Fleegler EW. Coronavirus Disease 2019 (COVID-19) and Firearms in the United States: Will an Epidemic of Suicide Follow? *Ann Intern Med* 2020; DOI: 10.7326/M20-1678.
135. COVID-19 Is Making America's Loneliness Epidemic Even Worse. Online document at: <https://time.com/5833681/loneliness-covid-19/>, accessed June 8, 2020.
136. Franceschi C, Garagnani P, Parini P, et al. Inflammaging: A new immune-metabolic viewpoint for age-related diseases. *Nat Rev Endocrinol* 2018;14:576–590.
137. Legido-Quigley H, Asgari N, Teo YY, et al. Are high-performing health systems resilient against the COVID-19 epidemic? *Lancet* 2020;395:848–850.
138. Finch CE, Crimmins EM. Inflammatory exposure and historical changes in human life-spans. *Science* 2004;305:1736–1739.
139. Peng C, Luttmann-Gibson H, Zanobetti A, et al. Air pollution influences on exhaled nitric oxide among people with type II diabetes. *Air Qual Atmos Health* 2016;9:265–273.
140. Abramson MJ, Wigmann C, Altug H, Schikowski T. Ambient air pollution is associated with airway inflammation in older women: A nested cross-sectional analysis. *BMJ Open Respir Res* 2020;7:e000549.
141. Finch CE, Kulminski AM. The Alzheimer's disease exposome. *Alzheimers Dement* 2019;15:1123–1132.
142. Moir RD, Lathe R, Tanzi RE. The antimicrobial protection hypothesis of Alzheimer's disease. *Alzheimers Dement* 2018;14:1602–1614.
143. Mazumder B, Almond D, Park K, et al. Lingering prenatal effects of the 1918 influenza pandemic on cardiovascular disease. *J Dev Orig Health Dis* 2010;1:26–34.
144. Dickson DA, Paulus JK, Mensah V, et al. Reduced levels of miRNAs 449 and 34 in sperm of mice and men exposed to early life stress. *Transl Psychiatry* 2018;8:101.
145. Cunningham AM, Walker DM, Nestler EJ. Paternal transgenerational epigenetic mechanisms mediating stress phenotypes of offspring. *Eur J Neurosci* 2019 [Epub ahead of print]; DOI: 10.1111/ejn.14582.

Address correspondence to:

William Bushell, PhD
Massachusetts Institute of Technology
Cambridge, MA 02139
 USA

E-mail: williamcbushell@gmail.com